Mechanistic evidence for the remote π -aryl
participation in acidcatalyzed ring opening of
homobenzoquinone epoxides

著者	Oshima Takumi, Asahara Haruyasu, Koizumi Takuya, Miyamoto Saki
journal or publication title	Chemical Communications
volume	15
page range	1804-1806
year	2008
URL	http://hdl.handle.net/10173/1352

doi: 10.1039/B719663F

Mechanistic evidence for the remote π -aryl participation in acid-catalyzed ring opening of homobenzoquinone epoxides

Takumi Oshima*, Haruyasu Asahara, Takuya Koizumi and Saki Miyamoto

Received (in XXX, XXX) 1st January 2007, Accepted 1st January 2007

First published on the web 1st January 2007

DOI: 10.1039/b000000x

The acid-induced reaction of bis(*p*-chlorophenyl)-homobenzoquinone epoxide gave the dual *ipso/ortho* intramolecular S_E2 -Ar products associated with the π -aryl participated oxirane ring opening, whereas bis(*p*-tolyl)- and diphenyl-substituted homologues provided only the *ortho* products.

The π -aryl participation is one of the most important physicochemical phenomena which control the reactivity of substrates and govern the reaction mechanism.¹ Such effects are generally ascribed to (derived from) the through-space electronic stabilization of the transition states by the direct electronic donation (not by resonance) of π -electrons from the aryl groups to the incipient carbocation center.² For instance a large number of studies have been made of the π -aryl assisted solvolyses of β -aryltosylates and brosylates from the kinetic³ and stereochemical⁴ point of view. By contrast, a little is known for the remote anchimeric assistance of aryl groups located in the carbon linkage far away from the reaction site.⁵ Thus, the elucidation of the possible remote π -aryl participation provides a further useful insight into the mechanistic understanding of the reactions involving the through-space π -electronic interaction.

Very recently, we found that the BF_3 -catalyzed ring-opening of diphenylhomobenzoquinone epoxide **1b** resulted in the transannular S_E2 -Ar displacement at *o*-position to afford tricyclic diketo-alcohol **3b** (Scheme 1).⁶ This reaction becomes of interest in that the endo-aromatic ring is likely to display the remote π -aryl participation in oxirane ring opening. Therefore, we felt that an appropriately *p*-substituted diphenylhomobenzoquinone epoxide **1** might allow to provide a possible *ipso*-product from the π -aryl participated transition state. Herein, we wish to report the mechanistic evidence for the very rare π -aryl-assisted oxirane ring opening in the BF_3 -catalyzed reaction of bis(*p*-chlorophenyl)homobenzoquinone epoxide **1c**.

The acid-induced reaction of *p,p'*-dimethyl-, unsubstituted, and *p,p'*-dichloro-substituted **1a-c** (0.02 mmol) was carried out in the presence of BF_3 (0.40 mmol) in $CDCl_3$ (0.62 ml) at ordinary temperature.† The reaction proceeded in a regioselective oxirane ring-opening at the Me substituted C-O bond and on treatment with water gave the common *o*-phenylene bridged tricyclic diketo-alcohols **3a-c** (for **3c** (20%)), as a mixture of its epimer **4c** (25%) and 2,5-cyclohexadien-4-one spiro-linked tricyclic diketo-alcohol **2c** (47%) for only the chloro-substituted **1c** in almost quantitative total yields based on the consumed **1** (Scheme 1).

Scheme 1 A dual pathway in BF_3 -catalyzed rearrangement of **1**.

Fig. 1 ORTEP representation (50% ellipsoids) of the structure **2c**.

The structures of new compounds **2c**, **3a**, **3c**, and **4c** were

deduced from their ^1H - and ^{13}C -NMR spectra and the **2c** was also confirmed by the X-ray crystal analysis (Fig. 1). §

As shown in Scheme 1, the formation of **2c** and **3a-c** can be rationalized by the occurrence of the competitive *ipso*- and *ortho*- $\text{SE}_2\text{-Ar}$ reaction via aryl bridged benzenonium ions, i.e., σ -complexes I and II (path a and path b), respectively. Although the *ortho*-bound intermediate II easily undergoes a rearomatization to afford **3a-c** via a proton migration, the formation of compound **2c** can be explained by the capture of the *ipso*-bound intermediate I with some water followed by the loss of HCl. Thus, the isolation of both the **2c** and **3c** can be taken as a strong evidence for the intervention of two σ -complexes, I and II. These schematic considerations prompted us to further examine the following mechanistic questions about the transition state leading to these σ -complexes⁷ as well as the marked substituent effects on the product distributions.

(1) Which can better explain the initial oxirane ring-opening, a concerted $\text{S}_{\text{N}}2$ -like pathway involving a π -aryl-assisted transition state or a stepwise $\text{S}_{\text{N}}1$ -like pathway generating a tertiary carbocation intermediate?

(2) Why does the *p*-chloro-substituted **1c** provide the dual *ipso/ortho* conjunct products in contrast to the *p,p'*-dimethyl-substituted **1a** and the unsubstituted **1b**?

As to the first question, the kinetic solvent effects provide a useful mechanistic information on the transition state. Namely, the more polar solvent will stabilize the polar transition state and largely accelerate the rate like in the $\text{S}_{\text{N}}1$ reactions.⁸ We have measured the rate constants for the MeSO_3H -catalyzed oxirane ring-opening of the parent unsubstituted epoxide **1b** by monitoring its first-order decay in various less basic solvents (Fig. 2). ¶ This reaction also gave the same tricyclic diketo-alcohol **3b** in almost quantitative yield as the BF_3 -catalyzed reaction. The observed

Fig. 2 A representative time course of the MeSO_3H ([30 mM])-catalyzed rearrangement of **1b** into **3b** in CDCl_3 (650 μl) at 30°C .

Table 1 Rate constants for MeSO_3H -catalyzed ring-opening of epoxide **1b** in various solvents at 30°C

Solvent	$E_{\text{T}}(30)$	k_2^a ($10^3, \text{M}^{-1}\text{s}^{-1}$)	k_{rel}
1,2-Dichloroethane	41.3	1.15	3.0
Dichloromethane	40.7	1.17	3.1
Chloroform- <i>d</i>	39.0	0.979	2.6
<i>o</i> -Dichlorobenzene	38.0	0.280	0.73
Fluorobenzene	37.0	0.380	0.99
Chlorobenzene	36.8	0.297	0.77
Benzene	34.3	0.384	1.0

^a The second-order rate constants k_2 were obtained by dividing the pseudo-first-order rate constants k_{obs} by the catalyst concentration ([30 mM]).

rate constants in a wide range of solvents at 30°C are summarized along with the solvent polarity parameter $E_{\text{T}}(30)$ ⁹ (Table 1). The total variation of k_2 amounts to only a factor of 3 over a wide range of solvent polarities investigated. The very poor kinetic solvent effects strongly support a concerted mechanism involving a less polar transition state. This observation is consistent with the appearance of the transition state in which the charge is highly dispersed on the π -aryl participated aromatic nucleus as well as on the breaking oxirane carbon atom.¹⁰ In such a $\text{S}_{\text{N}}2$ -like transition state, it is conceived that the orbital interaction between the HOMO of the π -electron donating aromatic group and the Walsh-type LUMO of oxirane ring¹¹ plays a crucial role in the cleavage of the relevant C-O bond as depicted in Scheme 1. The aryl participation in the ring opening of oxiranes is scarcely reported but has been put forwarded in order to explain the *syn*-stereochemistry in the acid-induced ring opening of a particular case of oxiranes bearing aryl groups directly or indirectly linked to the epoxide ring such as stilbene oxides¹² and spiro-linked 2-phenyl-1,2-epoxide¹³ or 1-benzyl-1,2-epoxides¹⁴ in which the well-documented phenonium ion intermediates are invoked.

The second question can be easily solved by considering the characteristic electronic properties of *p*-Cl substituent as exhibiting the electron-donating resonance effect as well as the good leaving ability which would stabilize the adjoining positive center of I and then enhance the release of HCl (Scheme 1). As to the *ipso*-attack, the *p*-tolyl and phenyl groups would rather facilitate such a reaction more efficiently than the *p*-chlorophenyl group. However, even if formed, such *ipso* σ -complexes of **1a** and **1b** would be inevitably transformed into the *ortho* σ -complex via a facile 1,2-shift because of the lack of the leaving ability of *p*-Me group (and of *p*-H atom). As a result, the lability of *ipso* intermediate I of **1c** toward residual water play a decisive role in the present product partitioning steps from the common transition state (Scheme 1).

In summary, we have succeeded in isolating both the *ipso*- and *ortho*- $\text{SE}_2\text{-Ar}$ products in the acid-catalyzed reaction of bis(*p*-chlorophenyl)-substituted homobenzoquinone epoxide **1c**. The present dual pathway for **1c** as well as the kinetic solvent effects is likely to prove that the acid-catalyzed ring-opening of diarylhomobenzoquinone epoxides **1** occurs via a concerted manner involving a very rare remote (δ -located) π -aryl participated transition state. The information obtained in the present reactions will provide a

useful insight into the understanding of Lewis acid-induced rearrangements of polycyclic epoxides.

H.A. expresses his special thanks for The Global COE (center of excellence) program “Global Education and Research Center for Bio-Environmental Chemistry” of Osaka University.

Notes and references

Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Yamada-oka, Suita, Osaka, 565-0871, Japan. E-mail: oshima@chem.eng.osaka-u.ac.jp

† Electronic Supplementary Information (ESI) available: Characterization data for the new substrates, **1a** and **1c**, and the products, **3a**, **3c**, **4c**. See DOI: 10.1039/b000000x/

‡ Representative procedure for acid-catalyzed rearrangement: To a solution of **1c** (0.02 mmol, 7.75mg) in 0.62 ml of CDCl₃ was added BF₃·OEt₂ (0.40 mmol, 50.2 μl). After standing for requisite time at ordinary temperature, the reaction mixture was quenched by water (5 ml) and extracted with CHCl₃ (5 ml × 3). The combined organic extracts were dried over anhydrous MgSO₄ and evaporated under reduced pressure. The residual mixture was submitted for ¹H NMR measurement for the determination of conversion of **1c** as well as the yields of **2c** and **3c**(**4c**). The reaction mixture was then purified by column chromatography on silica gel to successively afford **2c** and **3c** (as a mixture with **4c**) with hexane-benzene as eluent. The pure **4c** was obtained on treatment of **3c** with a few drops of Et₃N in CDCl₃ (0.6 ml) for 24h. The conversions of **1a**, **1b**, and **1c** were 100% (for 0.5h), >99 (4h), and 82 (20h), respectively. § The compound of **2c** has the following analytical data: mp 206.5–207 °C, colorless prisms (from hexane-chloroform). ¹H NMR (CDCl₃, 270 MHz, ppm): δ 1.00 (s, 3H), 1.08 (s, 3H), 2.75 (s, 1H), 2.93 (s, 1H), 4.00 (s, 1H), 6.17 (dd, *J* = 1.81, 10.4 Hz, 1H), 6.52 (dd, *J* = 1.81, 10.2 Hz, 1H), 6.54 (dd, *J* = 3.13, 10.4 Hz, 1H), 6.82 (dd, *J* = 3.13, 10.2 Hz, 1H), 7.00–7.10 (m, 2H), 7.25–7.26 (m, 2H). ¹³C NMR (CDCl₃, 75 MHz, ppm): δ 10.9, 14.8, 29.8, 43.1, 46.0, 52.8, 56.0, 75.4, 128.9, 129.7, 130.5, 131.3, 134.4, 135.3, 142.7, 147.5, 184.0, 203.0, 204.0. IR (KBr): 3417, 2925, 1745, 1664, 1261, 1091, 801 cm^{−1}.

Crystal data. **2c**: C₂₁H₁₇O₄Cl, *M* = 368.82, monoclinic, *a* = 11.4880(7), *b* = 12.5251(10), *c* = 13.3085(6) Å, β = 114.312(1)°, *V* = 1745.1(2) Å³, *T* = 23.0 °C, space group P2₁/n (#14), *Z* = 4, μ(MoKα) = 2.43 cm^{−1}, 14930 reflections measured, 3986 were unique (*R*_{int} = 0.070), *R*1[*I* > 2.0σ(*I*)] = 0.0901, *wR*2 (all data) = 0.2083. CCDC 666903.

¶ Since BF₃ is very sensitive to the residual water in the solvents employed, we investigated the kinetic solvent effects by using water-persistent MeSO₃H. The decay of **1b** was monitored by ¹H NMR for CDCl₃ and by HPLC for other solvents.

- (a) C. J. Lancelot, D. J. Cram and P. v. R. Scheyer, *Carbenium Ions*; G. A. Olah and P. v. R. Scheyer, Eds., Wiley-Interscience, New York, 1972, Vol. 3, Chapter 27, p 1347; (b) M. B. Smith and J. March, *March's Advanced Organic Chemistry*, Wiley, New York, 2001, Chapter 10, p 407; (c) M. Peeran, J. W. Wilt, R. Subramanian and D. S. Crumrine, *J. Org. Chem.*, 1993, **58**, 202; (d) D. N. Kevill and M. J. D'Souza, *J. Chem. Soc. Perkin Trans. 2*, 1997, 257; (e) M. Fujio, N. Goto, T. Dairokuno, M. Goto, Y. Saeki, Y. Okusaka and Y. Tsuno, *Bull. Chem. Soc. Jpn.*, 1992, **65**, 3072; (f) S. Nagumo, M. Ono, Y. Kakimoto, T. Furukawa, T. Hisano, M. Mizukami, N. Kawahara and H. Akita, *J. Org. Chem.*, 2002, **67**, 6618; (g) E. d. Río, M. I. Menéndez, R. López and T. Sordo, *J. Am. Chem. Soc.*, 2001, **123**, 5064.
- N. S. Isaacs, *Physical Organic Chemistry*, Longman Science & Technical, Essex, 1995, Chapter 13, p 643.
- (a) M. Fujio, K. Funatsu, M. Goto, M. Mishima and Y. Tsuno, *Tetrahedron*, 1987, **43**, 307; (b) M. Fujio, M. Goto, M. Mishima and Y. Tsuno, *Bull. Chem. Soc. Jpn.*, 1990, **63**, 1121; (c) M. Okamura, K. Hazama, M. Ohta, K. Kato and T. Horiguchi, *Chem. Lett.*, 1997, **26**, 973; (d) Y. Tsuno and M. Fujio, *Advances in Physical Organic Chemistry*; D. Bethell, Ed., Academic Press, London, 1999, Vol 32, p 267.

- (a) D. J. Cram, *J. Am. Chem. Soc.*, 1952, **74**, 2129, 2137; (b) A. Jr. Streitwieser, T. D. Walsh and J. R. Wolfe, *J. Am. Chem. Soc.*, 1965, **87**, 3686; (c) D. J. Cram and J. A. Thompson, *J. Am. Chem. Soc.*, 1967, **89**, 6766; (d) J. A. Thompson and D. J. Cram, *J. Am. Chem. Soc.*, 1969, **91**, 1778.
- (a) L. M. Jackman and V. R. Haddon, *J. Am. Chem. Soc.*, 1974, **96**, 5130; (b) M. Gates, D. L. Frank and W. C. von Felten, *J. Am. Chem. Soc.*, 1974, **96**, 5138; (c) T. Ando; J. Yamawaki and Y. Saito, *Bull. Chem. Soc. Jpn.*, 1978, **51**, 219.
- H. Asahara, E. Kubo, K. Togaya, T. Koizumi, E. Mochizuki and T. Oshima, *Org. Lett.*, 2007, **9**, 3421.
- Structure and dynamics of σ- and π-complexes are described in detail for the electrophilic aromatic substitutions, see S. M. Hubig and J. K. Kochi, *J. Org. Chem.*, 2000, **65**, 6807.
- C. Reichardt, “*Solvents and Solvent Effects in Organic Chemistry*” Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, 2003, ch.5, pp147–328.
- C. Reichardt, *Chem. Rev.*, 1994, **94**, 2319.
- We have previously reported a similar type of less polar transition state in the acid-catalyzed intramolecular Se₂-Ar reaction of cyclobutene-fused diarylhomobenzoquinones, see K. Koizumi, K. Harada, E. Mochizuki, K. Kokubo and T. Oshima, *Org. Lett.*, **2004**, **6**, 4081.
- (b) P. D. Mollere and K. N. Houk, *J. Am. Chem. Soc.*, 1977, **99**, 3226; (c) H. Iwamura, T. Sugawara, Y. Kawada, K. Tori, R. Muneyuki and R. Noyori, *Tetrahedron Lett.*, 1979, **20**, 3449; (d) R. D. Bach and G. J. Wolber, *J. Am. Chem. Soc.*, 1984, **106**, 1410.
- R. E. Parker and N. S. Isaacs, *Chem. Rev.*, 1959, **59**, 737.
- Y. Kita, A. Furukawa, J. Futamura, K. Higuchi, K. Ueda and H. Fujioka, *Tetrahedron*, 2001, **57**, 815.
- (a) P. Costantino, P. Crotti, M. Ferretti and F. Macchia, *J. Org. Chem.*, 1982, **47**, 2917; (b) P. Crotti, M. Ferretti, F. Macchia and A. Stoppioni, *J. Org. Chem.*, 1984, **49**, 4706; (c) P. Crotti, M. Ferretti, F. Macchia and A. Stoppioni, *J. Org. Chem.*, 1986, **51**, 2759.