

Tailor-Made Synthesis of N,N,2,6-Tetrasubstituted 4-Nitroanilines by Three-Component Ring Transformation of Dinitropyridone

| | |
|------------------------------|---|
| 著者 | Le Song Thi, Asahara Haruyasu, Nishiwaki Nagatoshi |
| journal or publication title | European Journal of Organic Chemistry |
| volume | 2015 |
| number | 6 |
| page range | 1203-1206 |
| year | 2015-02 |
| URL | http://hdl.handle.net/10173/1470 |

doi: 10.1002/ejoc.201403652

Tailor-made Synthesis of *N,N,2,6*-Tetrasubstituted 4-Nitroanilines by Three Component Ring Transformation of Dinitropyridone


Song Thi Le,^[a] Haruyasu Asahara,^[a,b] and Nagatoshi Nishiwaki^{[a,b]*}

Abstract: The ring transformation of dinitropyridone afforded various kinds of 2,6-disubstituted-4-nitroanilines upon treatment with aliphatic ketones in the presence of ammonium acetate as a nitrogen source, wherein dinitropyridone behaved as the synthetic equivalent of unstable nitromalaldehyde. The benzene ring as well as the amino group of the nitroaniline framework was easily modified by only changing a ketone and the nitrogen source, which afforded *N,N,2,6*-tetrasubstituted 4-nitroanilines in good to excellent yields.


Ring transformation reactions provide a unique method for the synthesis of functionalized heterocyclic compounds, which are not easily prepared by other methods.^[1,2] We have reported an alternative method for the synthesis of nitropyridines **3** by the three component ring transformation (TCRT) of dinitropyridone **1** with aromatic ketones **2** in the presence of ammonium acetate (Scheme 1),^[2] wherein dinitropyridone **1** serves as the synthetic equivalent of unstable nitromalaldehyde.^[3] This reaction is initiated by the nucleophilic attack of the enol form of **2** followed by conversion to enamine **4**. When the amino group of enamine **4** attacks the 6-position (route a), 2-arylated 5-nitropyridines **3** are formed via bicyclic intermediates **5** (Scheme 1).^[4] However, the β -carbon of enamine **4** cannot attack the 6-position (route b) because it would form a sterically strained four membered ring. On considering this reaction mechanism, we predicted that another ring transformation would occur when aliphatic ketones **6** having two α -hydrogens, viz. α - and α' -, are employed instead of aromatic ketones **2** (Scheme 2). In this ring transformation, both the amino group and the β -carbon can attack the 6-position in the case of enamine **9**, which leads to the formation of nitropyridines **7** (route a) or nitroanilines **8** (route c).

2,6-Disubstituted 4-nitroanilines **8** are useful synthetic intermediates for functional materials such as inhibitors of cholesterol acyl transferase,^[5] and π -conjugated polymers^[6] and are key intermediates for the synthesis of compounds with antimicrobial activities^[7] and the β -diketiminato ligand.^[8] Moreover, the push-pull electronic property is crucial for developing organic materials with potential for applications in nonlinear optics. Generally, 2,6-disubstituted 4-nitroanilines **8** are prepared from the corresponding anilines by nitration under harsh conditions, wherein protection and deprotection of the

amino group are necessary.^[9] However, the preparation of 2,6-disubstituted anilines is restricted because of the following limitations of the Friedel-Crafts alkylation:^[10] 1) the monoalkylated product undergoes further alkylation to afford polyalkylated products, 2) it is difficult to introduce two different alkyl groups, 3) primary alkyl groups longer than the ethyl group cannot be introduced, 4) phenyl and vinyl groups cannot be introduced, and 5) aminated and nitrated benzenes do not facilitate the alkylation. We believe that our TCRT method will overcome these disadvantages of the Friedel-Crafts alkylation and facilitate the synthesis of 2,6-disubstituted 4-nitroanilines **8** by only changing ketones **6**.



Scheme 1. TCRT of dinitropyridone **1** with aromatic ketones **2** in the presence of ammonium acetate, leading to the formation of 6-arylated nitropyridines **3**.




Scheme 2. TCRT of dinitropyridone **1** with aliphatic ketones **6** in the presence of ammonium acetate, affording disubstituted nitropyridines **7** and nitroanilines **8**.

[a] Ms. T. S. Le, Dr. H. Asahara, Prof. Dr. N. Nishiwaki
School of Environment Science and Engineering
Kochi University of Technology
Tosayamada, Kami, Kochi 782-8502, Japan
E-mail: nishiwaki.nagatoshi@kochi-tech.ac.jp
<https://sites.google.com/site/kutnishiwakilab2/nishiwakilab>
[b] Dr. H. Asahara, Prof. Dr. N. Nishiwaki
Research Center for Material Science and Engineering
Kochi University of Technology

When dinitropyridone **1** was allowed to react, at 65 °C for 24 h, with 3-pentanone **6a** in the presence of 5 equiv. of ammonium acetate in ethanol, nitroaniline **8a**^[11] and nitropyridine **7a**^[11] were obtained in 50% and 44% yield, respectively, resulting from two kinds of TCRTs (Table 1, entry 1). When 10 equiv. of ammonium acetate were used, the ratio of **8a** to **7a** increased considerably without decreasing the total yield (entry 2); this indicates the presence of equilibrium between **10** and **11**, which are kinetically and thermodynamically controlled intermediates, respectively, that can interconvert via enamine **9**. In the present TCRT, the competitive thermal decomposition of ammonium acetate to ammonia, which was liberated from the reaction mixture as a gas, also occurred. When all the ammonium acetate was consumed, the TCRT could no longer proceed because of the lack of a nitrogen source. Thus, increasing the amount of ammonium acetate prolonged the actual reaction time, resulting in the predominance of nitroaniline **8a**. However, no additional change was observed on using larger amounts of ammonium acetate (entry 3). Also, it was found that heating for 24 h was necessary for the completion of the TCRT (entries 4-6).

Table 1. Optimization of reaction conditions for the TCRT




| Entry | NH ₄ OAc (Equiv.) | Time (h) | Yield (%) | |
|-------|------------------------------|----------|-----------|-----------|
| | | | 8a | 7a |
| 1 | 5 | 24 | 50 | 44 |
| 2 | 10 | 24 | 83 | 13 |
| 3 | 15 | 24 | 86 | 13 |
| 4 | 10 | 18 | 70 | 14 |
| 5 | 10 | 12 | 64 | 13 |
| 6 | 10 | 6 | 55 | 13 |

This TCRT was applied to other ketones **6b-i** under the conditions optimized for **6a** (Table 2). When acetone **6b** was used as a substrate, two kinds of TCRTs occurred, similar to **6a**, to afford nitroaniline **8b**^[11] and nitropyridine **7b**^[11] in 51% and 47% yields, respectively (entry 2). It was possible to modify the 2- and 6- positions of the nitroaniline framework by only changing ketone **6** (entries 3-9). Notably, this TCRT facilitates the introduction of a propyl or phenyl group into the benzene ring, which cannot be achieved by the Friedel–Crafts reaction. As a result, symmetrical and unsymmetrical nitroanilines **8f-i**^[11] were easily prepared; however the yield of **8i** was low, presumably because steric repulsion by the phenyl groups prevents the formation of **11i** (entries 6-9).

As mentioned above, two kinds of TCRTs competitively occurred to form 5-nitropyridines **7** and 4-nitroanilines **8**. In these reactions, ammonium acetate serves as both a nitrogen source and an activator of ketone **6**. We believe that a combination of amine **12** and acetic acid, used instead of ammonium acetate, can carry out these roles, thus achieving the modification of the benzene ring as well as the amino group of the nitroaniline framework. In this case, only nitroanilines **13** will be formed as a TCRT product, because the aromatization of the intermediate, which is required for the formation of nitropyridines, is prevented by the *N*-substituents (R³ and R⁴).


Table 2. Application of the TCRT to other aliphatic ketones **6**



| Entry | Ketone 6 | | Product | Yield (%) | |
|-------|-------------------------------|-------------------------------|----------|--------------------------|--------------------------|
| | R ¹ | R ² | | 8 ^[11] | 7 ^[11] |
| 1 | Me | Me | a | 83 | 13 |
| 2 | H | H | b | 51 | 47 |
| 3 | Et | H | c | 66 | 10 |
| 4 | <i>i</i> -Pr | H | d | 58 | 0 |
| 5 | Pr | H | e | 83 | 9 |
| 6 | Et | Et | f | 67 | 24 |
| 7 | Pr | Pr | g | 74 | 22 |
| 8 | C ₆ H ₅ | Pr | h | 62 | 24 |
| 9 | C ₆ H ₅ | C ₆ H ₅ | i | 8 | 81 |

Propylamine **12A** was added to a solution of dinitropyridone **1**, 3-pentanone (**6a**) and acetic acid in ethanol, and the resulting solution was heated at 65 °C for 24 h. After the usual work-up, 2,6-dimethyl-4-nitro-*N*-propylaniline (**13Aa**) was obtained in 99% yield (Table 3, entry 1). This method was applied to the secondary amines, pyrrolidine **12B** and diethylamine **12C**, to afford *N,N*,2,6-tetrasubstituted 4-nitroanilines **13Ba** and **13Ca**, respectively, in excellent yields (entries 2 and 3). Methyl ketones **6b-d** also underwent this TCRT using a combination of either propylamine **12A** or pyrrolidine **12B** with acetic acid to afford the corresponding nitroanilines **13** in moderate to excellent yields (entries 4-9). Moreover, these reactions could induce modifications at the 2- and 6-positions using ketones **6e-h**, by which a propyl or a phenyl group could be introduced to the nitroaniline framework (entries 10-15).

Table 3. TCRT of dinitropyridone **1** with aliphatic ketones **6** using the mixture of amine **12** and acetic acid.



| Entry | Ketone 6 | | Amine 12 | | Product | Yield (%) |
|-------|-------------------------------|-------------------------------|------------------------------------|----------------|----------|----------------|
| | R ¹ | R ² | R ³ | R ⁴ | | |
| 1 | Me | Me | Pr | H | A | 13Aa 99 |
| 2 | Me | Me | -(CH ₂) ₄ - | H | B | 13Ba 98 |
| 3 | Me | Me | Et | Et | C | 13Ca 98 |
| 4 | Et | H | Pr | H | A | 13Ab 83 |
| 5 | Et | H | -(CH ₂) ₄ - | H | B | 13Bb 68 |
| 6 | Pr | H | Pr | H | A | 13Ac 77 |
| 7 | Pr | H | -(CH ₂) ₄ - | H | B | 13Bc 87 |
| 8 | Pr | H | Et | Et | C | 13Cc 51 |
| 9 | <i>i</i> -Pr | H | Pr | H | A | 13Ad 83 |
| 10 | Et | Et | Pr | H | A | 13Ae 69 |
| 11 | Et | Et | -(CH ₂) ₄ - | H | B | 13Be 68 |
| 12 | Pr | Pr | Pr | H | A | 13Af 81 |
| 13 | Pr | Pr | -(CH ₂) ₄ - | H | B | 13Bf 59 |
| 14 | C ₆ H ₅ | Pr | Pr | H | A | 13Ag 80 |
| 15 | C ₆ H ₅ | C ₆ H ₅ | Pr | H | A | 13Ah 32 |

In summary, we have developed a new preparative method for 2,6-disubstituted 4-nitroanilines **8** and **13** by the TCRT of dinitropyridone **1** with aliphatic ketones **6** in the presence of ammonium acetates. In this reaction, a number of ketones **6** are usable as substrates, which facilitate the modification of the nitroaniline framework. In addition, this TCRT requires only simple experimental manipulations and mild reaction conditions, which is advantageous from the viewpoint of practical use. These features facilitate the construction of a library of compounds that are not easily available by other methods. Furthermore, modification of the amino group was successfully achieved by using a combination of amine **12** and acetic acid. Consequently, the tailor-made synthesis of *N,N*,2,6-tetrasubstituted 4-nitroanilines **8** and **13** became possible on demand.

Experimental Section

To a solution of the nitropyridone **1** (50 mg, 0.25 mmol) in ethanol (5 mL), were added 3-pentanone (**6a**, 26 μ L, 0.25 mmol) and ammonium acetate (96.3 mg, 1.25 mmol), and then the resultant mixture was heated at 65 °C for 24 h. After removal of the solvent, the residue was washed with benzene (3 \times 10 mL) to remove unreacted ketone **6a**, which affords a mixture of the nitropyridine **7a** and the nitroaniline **8a**. The separation of products was performed by column chromatography on silica gel (hexane : ethyl acetate = 95 : 5) to afford **7a** (18.3 mg, 0.11 mmol, 44%) and **8a** (20.8 mg, 0.13 mmol, 50%), respectively. The TCRT reaction of the dinitropyridone **1** with other ketones were performed in a similar way.

Acknowledgements

We sincerely thank Prof. Satoshi Minakata (Osaka University, Japan) for his kind assistance. This work was supported by JSPS KAKENHI Grant Number 26410123.

Keywords: Nitroheterocycles • Multicomponent reaction • Ring transformation • Ketones • Synthetic method.

- [1] **Diels-Alder type ring transformation:** a) J. D. Kirkham, R. J. Butlin, J. P. A. Harrity, *Angew. Chem. Int. Ed.* **2012**, *51*, 6402; b) C. Sabot, E. Oueis, X. Brune, P. Y. Renard, *Chem. Commun.* **2012**, *48*, 768; c) E. D. Anderson, D. L. Boger, *Org. Lett.* **2011**, *13*, 2492; d) T. Delaunay, P. Genix, M. Es-Sayed, J. P. Vors, N. Monterio, G. Balme, *Org. Lett.* **2010**, *12*, 3328; e) C. Wu, Y. Fang, R. C. Larock, F. Shi, *Org. Lett.* **2010**, *12*, 2234; f) T. Miura, M. Yamauchi, M. Murakami, *Chem. Commun.* **2009**, 1470; g) Y. Yoshino, T. Kurahashi, S.-J. Matsubara, *J. Am. Chem. Soc.* **2009**, *131*, 7494; h) H. Xie, L. Zu, H. R. Oueis, H. Li, J. Wang, W. Wang, *Org. Lett.* **2008**, *10*, 1923.
- ANROR type ring transformation:** a) H. G. Bonaccorso, J. Navarini, L. M. F. Porte, E. P. Pittaluga, A. F. Junges, A. R. Mayer, M. A. P. Martins, N. Zanatta, *J. Fluor. Chem.* **2013**, *151*, 38; b) P. A. Koutentis, M. Koyioni, S. S. Michaelidou, *Org. Biomol. Chem.* **2013**, *11*, 621; c) A. Rykowski, E. Wolinska, D. Branowska, H. C. van der Plas, *ARKIVOC* **2004**, *iii*, 74; d) G. Hajós, Z. Riedl, G. Kollenz, *Eur. J. Org. Chem.* **2001**, 3405; e) H. C. van der Plas, *J. Heterocycl. Chem.* **2000**, *37*, 427.
- Nucleophilic type ring transformation:** a) C. Henry, A. Haupt, S. C. Turner, *J. Org. Chem.* **2009**, *74*, 1932; b) G. P. Sagitullina, A. K. Garkushenko, Y. O. Vinokurova, V. A. Nyrkova, E. G. Atavin, R. S. Sagitullin, *Russ. Org. Chem.* **2009**, *45*, 1045; (c) N. Nishiwaki, M. Ariga, *Topics in Heterocyclic Chemistry Vol. 8* ed. by S. Eguchi, Springer, Berlin, **2007**, pp 43-72.
- [2] a) S. T. Le, H. Asahara, K. Kobi, R. Sugimoto, K. Saigo, N. Nishiwaki, *Asian. J. Org. Chem.* **2014**, *3*, 297; b) S. T. Le, H. Asahara, N. Nishiwaki, *Synthesis* **2014**, 2175.
- [3] N. Nishiwaki, S. Hirao, J. Sawayama, K. Saigo, *Heterocycles* **2012**, *84*, 115.
- [4] Similar reaction mechanism was proposed for the ring transformation of the nitropyrimidinone: N. Nishiwaki, R. Sugimoto, K. Saigo, K. Kobi, *Tetrahedron. Lett.* **2013**, *54*, 956.
- [5] H. Saka, M. Muraoka, S. Onuma, *Jpn. Kokai Tokkyo Koho*, **2002**-173476.
- [6] F. Chimenti, A. Boasco, D. Secci, P. Chimenti, A. Granese, *Synth. Commun.* **2004**, *34*, 2549.
- [7] C. M. Jamkhandi, J. I. Disouza, *J. Pharm. Sci.* **2013**, *5*, 225.
- [8] N. M. Rajendran, A. Haleel, N. Reddy, N. Dastagiri, *Organometallics* **2014**, *33*, 217.
- [9] a) F. J. Carver, C. A. Hunter, D. J. Livingstone, J. F. McCabe, E. M. Seward, *Chem.-Eur. J.* **2002**, *8*, 2848; b) S. Al-Khafaji, N. Cardinale, J. R. Hanson, *J. Chem. Res. Synop.* **2003**, 388, 701.
- [10] a) P. Vollhardt, K. P. C. Vollhardt, N. E. Schore, *Organic Chemistry Structure and Function*, 5th ed. New York: W. H. Freeman and Company, **2007**; b) L. G. Wade Jr., *Organic Chemistry*. 6th ed. New Jersey: Pearson Prentice Hall, **2006**.
- [11] For known compounds **7** and **8**; **7a**: Y. Tohda, M. Eiraku, T. Nakagawa, Y. Usami, M. Ariga, T. Kawashima, K. Tani, H. Watanabe, Y. Mori, *Bull. Chem. Soc. Jpn.* **1990**, *63*, 2820; **7b**: Y. Liu, P. Ren, K. Jessen, X. Guo, C. Rommel, T. Wilson, E. Troy, *PCT Int. Appl. WO* **2014**-151147; **7c**: A. S. Jørgensen, P. Jacobsen, L. B. Christiansen, P. S. Bury, A. Kanstrup, S. M. Thorpe, L. Naerum, K. Wassermann, *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2383; **7d**, **7f**, **7g** and **8h**: N. Nishiwaki, H. Tatsumichi, M. Tamura, M. Ariga, *Lett. Org. Chem.* **2006**, *3*, 629; **7e**: W. Gruber, *Can. J. Chem.* **1953**, *31*, 1181; **7i**: P. Barczynski, H. C. van der Plas, *Reu. Trav. Chim. Pays-Bas*, **1978**, *97*, 256; **8a**: T. Nanjo, C. Tsukano, Y. Takemoto, *Org. Lett.* **2012**, *14*, 4270; **8b**: C. Matheis, K. Jouvin, L. J. Goossen, *Org. Lett.* **2014**, *16*, 5984; **8c**: C. W. L. Bevan, G. C. Bye, *J. Chem. Soc.* **1957**, 3194; **8d**: M. Birch, G. E. M. Sibley, D. Law, J. D. Oliver, *PCT Int. Appl. WO* **2009**-144473; **8e**: G. Baddeley, J. Kenner, *J. Chem. Soc.* **1935**, 303; **8f**: K. Minksztym, A. Jarczewski, *J. Mol. Struct.* **2004**, *691*, 203; **8i**: D. Meinhard, M. Wegner, G. Kipiani, A. Hearley, P. Reuter, S. Fischer, O. Marti, B. Rieger, *J. Am. Chem. Soc.* **2007**, *129*, 9182.

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

Text for Table of Contents

((Insert TOC Graphic here))


Author(s), Corresponding Author(s)*

Page No. – Page No.

Title

Layout 2:

COMMUNICATION



The ring transformation of dinitropyridone afforded various kinds of *N,N,2,6*-tetrasubstituted 4-nitroanilines, of which the benzene ring as well as the amino group was easily modified by only changing a ketone and an amine.

Song Thi Le, Haruyasu Asahara, and
Nagatoshi Nishiwaki*

Page No. – Page No.

**Tailor-made Synthesis of *N,N,2,6*-
Tetrasubstituted 4-Nitroanilines by
Three Component Ring
Transformation of Dinitropyridone**