

An Efficient Synthesis of Nitrated Cycloalka[b]pyridines

著者	Le Song Thi, Asahara Haruyasu, Nishiwaki Nagatoshi
journal or publication title	Synthesis
volume	46
number	16
page range	2175-2178
year	2014
URL	http://hdl.handle.net/10173/1493

doi: 10.1055/s-0033-1341237

An Efficient Synthesis of Nitrated Cycloalka[*b*]pyridines

Thi Song Le, Haruyasu Asahara, and Nagatoshi Nishiwaki*

School of Environmental Science and Engineering, Kochi University of Technology, Tosayamada, Kami, Kochi 782-8502, Japan

Fax: +81-887-57-2520

E-mail: nishiwaki.nagatoshi@kochi-tech.ac.jp


Received: The date will be inserted once the manuscript is accepted.

Abstract: The three-component ring transformation of 1-methyl-3,5-dinitro-2-pyridone with cycloalkanones of different ring sizes in the presence of ammonium acetate afforded the corresponding nitrated cycloalka[*b*]pyridines in high yields. Furthermore, a double bond could be easily introduced to the product by changing the cycloalkanone to a cycloalkenone.

Keywords: pyridines, multicomponent reaction, ketones, heterocycles, bicyclic compounds, and ring transformation.

Nitro compounds are important organic compounds that are widely used in organic syntheses because of diverse reactivities.¹ We have reported a synthetic method for the arylated nitropyridines **3** by the three-component ring transformation (TCRT) of dinitropyridone **1** with aromatic ketones **2** in the presence of ammonia,² in which dinitropyridone **1** serves as the synthetic equivalent of unstable nitromalonaldehyde (Scheme 1, method a).³ Recently, other research groups also reported the TCRT of **1** with ketones using ammonia as the nitrogen source.⁴ Although these methods easily provide nitropyridine derivatives via a single step, they suffer from low yields of the products because of the competitive ammonolysis of the substrate **1**.⁵ Furthermore, methanolic ammonia should be prepared in advance, which is also troublesome.















These disadvantages were overcome by using less nucleophilic ammonium acetate (NH₄OAc) instead of ammonia.⁶ The reaction of dinitropyridone **1** with aromatic ketones **2** in the presence of NH₄OAc afforded the nitropyridines **3** in good to excellent yields (Scheme 1, method b).⁷ This TCRT is advantageous in terms of the use of solid NH₄OAc compared to gaseous ammonia, and extra NH₄OAc can be easily removed from the reaction mixture by thermal decomposition. The successful results of our study prompted us to extend the substrate scope of this method to a series of cyclic ketones **4**. Thus, nitrated cycloalka[*b*]pyridines **5** could be successfully obtained and further used as useful intermediates for the synthesis of metacyclophanes,⁴ pharmacophores,⁸ and biologically active compounds.⁹




Scheme 1. Synthetic methods for 2-arylated-5-nitropyridines **3** by TCRT of dinitropyridone **1** with aromatic ketones **2** in the presence of a nitrogen source.

The reaction of the dinitropyridone **1** with cyclohexanone **4a** in the presence of 5 equiv. NH₄OAc at 65 °C for 24 h afforded nitrated cyclohexa[*b*]pyridine **5a**^{2b,4b,10} in a moderate yield, because of the shortage of the nitrogen source due to the competitive thermal decomposition of NH₄OAc (Table 1, entry 1).⁷ This problem was easily solved by increasing the amount of NH₄OAc, and the yield of **5a** could be increased to 95% (entries 2 and 3). When microwave heating was used, nitropyridine **5a** was afforded efficiently within 1 h (entry 4). Cyclopentanone **4b** was less reactive than **4a**, thus affording cyclopentapyridine **5b**^{2b,10} in only 67% yield under the same reaction conditions used for **4a** (entry 5). In such a case, microwave heating was effective, and the reaction was completed within a short time under microwave heating (entries 6 and 7). In contrast, larger cycloalkanones **4c** and **4d** underwent the TCRT efficiently to afford the corresponding cyclohepta- and cyclooctapyridines **5c**¹⁰ and **5d**,^{4b,10} respectively (entries 8–11). When unsymmetrical 2-methylcyclohexanone **4e** was employed, 8-methylated tetrahydroquinoline **5e** was obtained efficiently (entries 12 and 13). This reaction was also applied to unsaturated cyclic ketone **4f**, thus affording 7,8-dihydroquinoline **5f** even though microwave heating was necessary for efficient TCRT (entries 14–16). On the other hand, cyclopentenone **4g** did not undergo the TCRT even though microwave heating was used (entry 17).

Table 1. Synthesis of cycloalka[*b*]pyridines **5** by TCRT.

Entry	n	Ketones	NH ₄ OAc (equiv.)	Time (h)	Product	Yield (%)	
1	2		4a	5		5a	56
2			10	24		88	
3			15	24		95	
4			15	1 ^a		97	
5	1		4b	15		5b	67
6			15	2 ^a		87	
7			15	1 ^a		70	
8	3		4c	15		5c	94
9			15	1 ^a		91	
10	4		4d	15		5d	85
11			15	1 ^a		95	
12	2		4e	15		5e	83
13			15	2 ^a		86	
14	2		4f	15		5f	59
15			15	3 ^a		89	
16			15	1 ^a		40	
17	1		4g	15		5g	0

a: Microwave heating was used.






Figure 1. Two plausible intermediates derived from 1-methylcyclohexanone **4e**.




Figure 2. The intermediate **7b** and **7f**.

A plausible mechanism for this TCRT is shown in Scheme 2. After the addition of the enol form of **4a** to the 4-position of **1**, the resulting adduct **6a** is transformed to enamine **7a** by reacting with ammonium ion. The intramolecular attack of the amino group at the 6-position affords tricyclic intermediate **8a**, from which nitroacetamide is eliminated to afford nitropyridine **5a** after the aromatization of intermediate **9a**. Although another reaction path via enamine **10a** can be considered, the same product **5a** is formed. In the case of unsymmetrical 2-methylcyclohexanone **4e**, two kinds of intermediates **8e** and **8e'** are possible (Figure 1), however, the latter intermediate cannot afford aromatized product. Therefore, only **5e** is formed via the intermediate **8e**. In the reactions using cyclopentanone **4b** and cyclohexenone **4f**, the loss of the flexibility of **7b** and **7f** makes the formation of tricyclic intermediates **8b** and **8f** more difficult (Figure

2). To the contrary, cyclopentenone **4g** did not afford ring transformed product **5g** because formation of a sterically restricted intermediate is necessary.

In summary, nitrated cycloalka[*b*]pyridines **5** were efficiently synthesized by the TCRT of dinitropyridone **1** with saturated and unsaturated cyclic ketones **4** in the presence of NH_4OAc . Because the method requires only simple manipulations and mild reaction conditions, it has great potential for the synthesis of [*b*]-condensed pyridine frameworks.

Experimental Section

The melting points were determined on a Yanaco micro-melting-points apparatus, and were uncorrected. The ^1H NMR spectra were measured on a Bruker Ascend-400 at 400 MHz with TMS as an internal standard. The ^{13}C NMR spectra were measured on a Bruker Ascend-400 at 100 MHz, and assignment was performed by DEPT experiments. The IR spectra were recorded on a JASCO FT/IR-4200 spectrometer. The high resolution mass spectrum was measured on a JEOL JMS-DX303HF. Microwave heating was performed using Anton-Paar Microwave 300. All the reagents and solvents were commercially available and were used as received.

General procedure of TCRT. To a solution of the dinitropyridone **1** (50 mg, 0.25 mmol) in ethanol (5 mL), cyclohexanone **4a** (26 μL , 0.25 mmol) and NH_4OAc (289 mg, 3.75 mmol) were added, and the resultant mixture was heated at 65 $^\circ\text{C}$ on the oil bath for 24 h. After removal of the solvent, the residue was washed with benzene (3 \times 10 mL) to afford 5,6,7,8-tetrahydro-3-nitroquinoline (**5a**) (42 mg, 0.24 mmol, 95%) as a yellow powder. The reactions of the dinitropyridone **1** with other ketones **4b-g** were performed in a similar way. When the reaction was conducted using microwave heating, the experiment was conducted in a similar way.

6,7-Dihydro-3-nitro-5*H*-cyclopenta[*b*]pyridine (**5b**).^{2b,10}

Yellow powder.

Yield: 43 mg, 97%.

6,7,8,9-Tetrahydro-3-nitro-5*H*-cyclohepta[*b*]pyridine (**5c**).¹⁰

Yellow powder.

Yield: 36 mg, 87%.

5,6,7,8,9,10-Hexahydro-3-nitro-cycloocta[*b*]pyridine (**5d**).^{4b,10}

Yellow powder.

Yield: 49 mg, 95%.

5,6,7,8-Tetrahydro-8-methyl-3-nitroquinoline (**5e**).¹⁰

Yellow powder.

Yield: 41 mg, 86%.

7,8-Dihydro-3-nitroquinoline **5f**.

Yellow powder; mp 57–59 $^\circ\text{C}$.

Yield: 40 mg, 89%.

IR (KBr, cm^{-1}) 1261, 1577, 1509.

^1H NMR (CDCl_3) δ = 2.49 (ddt, J = 2.8, 4.0, 8.4 Hz, 2H), 2.99 (t, J = 8.4 Hz, 2H), 6.57 (dt, J = 4.0, 8.0 Hz, 1H), 6.74 (ddt, J = 0.8, 2.8, 8.0 Hz, 1H), 8.16 (dd, J = 0.8, 2.8 Hz, 1H), 9.19 (d, J = 2.8 Hz, 1H).

^{13}C NMR (CDCl_3) δ = 22.6 (CH_2), 26.5 (CH_2), 128.9 (CH), 129.3 (CH), 131.3 (C), 139.1 (CH), 141.1 (C), 143.3 (CH), 158.8 (C).

HRMS (EI, magnetic field) Calcd for $\text{C}_9\text{H}_8\text{N}_2\text{O}_2$: 176.0586. Found: 176.0586.

Acknowledgment

We sincerely thank Prof. Satoshi Minakata (Osaka University, Japan) for his kind assistance.

References

- (1) (a) Nishiwaki, N. in *Comprehensive Organic Synthesis*, 2nd ed., Vol. 6, 100–130, Elsevier, Oxford, UK (2014); (b) Balini, R.; Gabrietti, S.; Palmieri, A.; Petrini, M. *Curr. Org. Chem.* **2011**, *15*, 1482–1506; (c) Ballini, R.; Barboni, L.; Fiorini, D.; Palmieri, A.; Petrini, M. *ARKIVOC* **2006**, vi, 127–152; (d) Adams, J. P.; Peterson, J. R. *J. Chem. Soc., Perkin Trans. 1* **2000**, 3695–3705.
- (2) Nishiwaki, N.; Ariga, M. in *Topics in Heterocyclic Chemistry, Bioactive Heterocycles II, Vol. 8*, Ed. by Eguchi, S., 43–72, Springer, Berlin (2007); (b) Tohda, Y.; Kawahara, T.; Eiraku, M.; Tani, K.; Nishiwaki, N.; Ariga, M. *Bull. Chem. Soc. Jpn.* **1994**, *67*, 2176–2186; (c) Tohda, Y.; Eiraku, M.; Nakagawa, T.; Usami, Y.; Ariga, M.; Kawashima, T.; Tani, K.; Wanatabe, H.; Mori, Y. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 2820–2827.
- (3) Nishiwaki, N.; Hirao, S.; Sawayama, J.; Saigo, K. *Heterocycles* **2012**, *84*, 115–134.
- (4) (a) Henry, C.; Haupt, A.; Turner, S. C. *J. Org. Chem.* **2009**, *74*, 1932–1938; (b) Sagitullina, G. P.; Garkushenko, A. K.; Vinokurova, Y. O.; Nyrkova, V. A.; Atavin, E. G.; Sagitullin, R. S. *Russ. J. Org. Chem.* **2009**, *45*, 1045–1049.
- (5) Tohda, Y.; Ariga, M.; Kawashima, T.; Matsumura, E. *Bull. Chem. Soc. Jpn.* **1987**, *60*, 201–204.
- (6) The TCRT of 3-methyl-5-nitro-4-pyrimidinone with ketone in the presence of NH_4OAc efficiently proceeds to afford pyrimidines, nitropyridones and aminopyridines. (a) Nishiwaki, N.; Sugimoto, R.; Saigo, K.; Kobihiro, K. *Tetrahedron Lett.* **2013**, *54*, 956–959; (b) Nishiwaki, N.; Yamashita, K.; Azuma, M.; Adachi, T.; Tamura, M.; Ariga, M. *Synthesis* **2004**, 1996–2000; (c) Nishiwaki, N.; Azuma, M.; Tamura, M.; Hori, K.; Tohda, Y.; Ariga, M. *Chem. Commun.* **2002**, 2170–2171.
- (7) Le, T. S.; Asahara, H.; Kobihiro, K.; Sugimoto, R.; Saigo, K.; Nishiwaki, N. *Asian. J. Org. Chem.* **2014**, *3*, 297–302.
- (8) Yamada, S.; Goto, T.; Yamaguchi, T.; Aihara, K.; Kogi, K.; Narita, S. *Chem. Pharm. Bull.* **1995**, *43*, 421–431.
- (9) Yamada, S.; Goto, T.; Shimanuki, E.; Narita, S. *Chem. Pharm. Bull.* **1994**, *42*, 718–720.
- (10) Marcellis, A. T. M.; van der Plas, H. C. *Tetrahedron* **1989**, *45*, 2693–2702.